



**You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice**

Title: Pentagon diagrams of Bhabha scattering

Author: J. Fleischer, Janusz Gluza, K. Kajda, T. Riemann

Citation style: Fleischer J., Gluza Janusz, Kajda K., Riemann T. (2007).
Pentagon diagrams of Bhabha scattering. "Acta Physica Polonica B" (Vol. 38,
no. 11 (2007), s. 3529-3536).



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie,
rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem
oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego

PENTAGON DIAGRAMS OF BHABHA SCATTERING* **

J. FLEISCHER^{a,b}, J. GLUZA^c, K. KAJDA^c, T. RIEMANN^b

^aFakultät für Physik, Universität Bielefeld
Universitätsstr. 25, 33615 Bielefeld, Germany

^bDeutsches Elektronen-Synchrotron DESY
Platanenalle 6, 15738 Zeuthen, Germany

^cInstitute of Physics, University of Silesia
Uniwersytecka 4, 40-007 Katowice, Poland

(Received October 29, 2007)

We report on tensor reduction of five point integrals needed for the evaluation of loop-by-loop corrections to Bhabha scattering. As an example we demonstrate the calculation of the rank two tensor integral with cancellation of the spurious Gram determinant in the denominator. The reduction scheme is worked out for arbitrary five point processes.

PACS numbers: 13.20.Ds, 13.66.De

1. Introduction

Evaluation of the Bhabha cross-section with two-loop accuracy [1–4] would be incomplete without taking into account contributions from one-loop diagrams with photon emission from internal electron lines. So far bremsstrahlung from external legs (lowest order) has been taken into account [5] and implemented in [6–8].

The general form of the five point tensor integrals in the diagrams of Fig. 1 is:

$$I_5^{\{1,q^\mu,q^\mu q^\nu \dots\}} = e^{\varepsilon\gamma_E} \int \frac{d^d q}{i\pi^{d/2}} \frac{\{1,q^\mu,q^\mu q^\nu \dots\}}{c_1 c_2 c_3 c_4 c_5}, \quad (1.1)$$

where $c_i = (q + q_i)^2 - m_i^2$, $i = 1, \dots, 5$. We also make the (arbitrary) choice $q_5 = 0$.

* Presented by K. Kajda at the XXXI International Conference of Theoretical Physics, “Matter to the Deepest”, Ustroń, Poland, September 5–11, 2007.

** Work supported in part by Sonderforschungsbereich/Transregio TRR 9 of DFG ‘Computergestützte Theoretische Teilchenphysik’, and by the European Community’s Marie Curie Research Training Networks MRTN-CT-2006-035505 ‘HEPTOOLS’ and MRTN-CT-2006-035482 ‘FLAVIANet’.




Fig. 1. Eight diagrams at one-loop level obtained using DIANA [11].

Massive five point functions tend to be unstable numerically in certain kinematic domains due to the appearance of an inverse Gram determinant. Calculating the tensor integrals directly, *e.g.* by using Mellin–Barnes (MB) representations, we have to cope with five dimensional MB integrals (before expansion in epsilon). After ϵ expansion, five dimensional MB representation still remains at ϵ^0 (*e.g.* for second order tensor). The MB representations are derived with the aid of the Mathematica packages **AMBRE** and **MB** [9, 10]. They are hard to evaluate both numerically (and in fact presently the MB package works in Euclidean region only) and analytically. A reduction of tensor five point functions to scalar four and three point functions solves that problem.

2. Reduction of five point functions

We give examples of reduction for the scalar and second rank tensor five point function where the spurious Gram determinant in denominators is avoided (see also [12]).

To make a reduction one can follow the standard Passarino–Veltman [13] reduction scheme, or, as we will do, follow a scheme based on [14–16]. For a diagram with internal lines $1 \dots n$ we can introduce the so called “Modified Cayley Determinant”:

$$(\cdot)_5 = \begin{vmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & Y_{11} & Y_{12} & \dots & Y_{1n} \\ 1 & Y_{12} & Y_{22} & \dots & Y_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & Y_{1n} & Y_{2n} & \dots & Y_{nn} \end{vmatrix}, \quad (2.1)$$

with $Y_{ij} = -(q_i - q_j)^2 + m_i^2 + m_j^2$. By cutting from $()_5$ rows j_1, j_2, \dots and columns k_1, k_2, \dots we get so-called “Minors”. The sign of a “Signed Minor”,

$$\begin{pmatrix} j_1 & j_2 & \dots \\ k_1 & k_2 & \dots \end{pmatrix}_n, \quad (2.2)$$

is determined by the sum of indices of excluded rows and columns and by taking into account the appropriate signatures of the permutations, taken separately from excluded rows and columns (important when cancelling more than one row and one column), *e.g.* signatures of permutations for $\begin{pmatrix} 12 \\ 54 \end{pmatrix}$ is: + for rows and – for columns. In this way:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}_n = \begin{vmatrix} Y_{11} & Y_{12} & \dots & Y_{1n} \\ Y_{12} & Y_{22} & \dots & Y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ Y_{1n} & Y_{2n} & \dots & Y_{nn} \end{vmatrix}. \quad (2.3)$$

Using “signed minors” one can write tensor reduction formulas in a compact and elegant way.

We start from the simplest case of a scalar five point function. As a starting point we use the recursion relation [15]:

$$(d - \sum_{i=1}^n \nu_i + 1) G_{n-1} I_n^{(d+2)} = \left[2\Delta_n + \sum_{k=1}^n \left(\frac{\partial \Delta_n}{\partial m_k^2} \right) \mathbf{k}^- \right] I_n^{(d)}, \quad (2.4)$$

where

$$\Delta_n = \begin{pmatrix} 0 \\ 0 \end{pmatrix}_n, \quad G_{n-1} = - \sum_{k=1}^n \left(\frac{\partial \Delta_n}{\partial m_k^2} \right) = 2()_n. \quad (2.5)$$

The operator \mathbf{k}^- decreases the k -th propagator power by one. Because in our case powers of propagators are all equal to one, and we are interested in the case with $n = 5$, the above relation changes into:

$$(d - 4)()_5 I_5^{(d+2)} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}_5 I_5 - \sum_{k=1}^5 \begin{pmatrix} 0 \\ k \end{pmatrix}_5 I_4^k. \quad (2.6)$$

In the limit $d \rightarrow 4$ we get as the final reduction formula for the scalar five point function:

$$I_5 = \frac{1}{\begin{pmatrix} 0 \\ 0 \end{pmatrix}_5} \sum_{k=1}^5 \begin{pmatrix} 0 \\ k \end{pmatrix}_5 I_4^k, \quad (2.7)$$

where I_4^k is the scalar four point integral obtained by cancelling the k -th line in the five point function diagram.

For the vector integral we refer to [15]. The second rank tensor integral can be written in the form:

$$I_5^{\mu\nu} = \sum_{i,j=1}^4 q_i^\mu q_j^\nu n_{ij} I_{5,ij}^{[d+]^2} - \frac{1}{2} g^{\mu\nu} I_5^{[d+]}. \quad (2.8)$$

Here $[d+]^l = 4 + 2l - 2\varepsilon$, and $n_{ij} = 1 + \delta_{ij}$. We start from the following formulas [15]:

$$n_{ij} I_{5,ij}^{[d+]^2} = -\frac{\binom{0}{j}_5}{\binom{0}{5}} I_{5,i}^{[d+]} + \sum_{s=1, s \neq i}^5 \frac{\binom{s}{j}_5}{\binom{s}{5}} I_{4,i}^{[d+],s} + \frac{\binom{i}{j}_5}{\binom{i}{5}} I_5^{[d+]}, \quad (2.9)$$

$$g^{\mu\nu} = 2 \sum_{i,j=1}^4 \frac{\binom{i}{j}_5}{\binom{i}{5}} q_i^\mu q_j^\nu, \quad (2.10)$$

$$I_{4,i}^{[d+],s} = -\frac{\binom{0s}{is}_5}{\binom{s}{5}} I_4^s + \sum_{t=1, t \neq s}^5 \frac{\binom{ts}{is}_5}{\binom{s}{5}} I_3^{st}. \quad (2.11)$$

Using all the above contributions we end up with:

$$I_5^{\mu\nu} = \sum_{i,j=1}^4 q_i^\mu q_j^\nu I_{5,ij}, \quad (2.12)$$

where

$$I_{5,ij} = \frac{1}{\binom{0}{5}} \left\{ -\frac{\binom{0}{j}_5}{\binom{0}{5}} \sum_{s=1}^5 \binom{0i}{0s}_5 I_4^s - \sum_{s=1}^5 \frac{\binom{s}{j}_5 \binom{0s}{is}_5}{\binom{s}{5}} I_4^s + \sum_{s,t=1}^5 \frac{\binom{s}{j}_5 \binom{ts}{is}_5}{\binom{s}{5}} I_3^{st} \right\}. \quad (2.13)$$

However, we can avoid the determinant $\binom{0}{5}$ in the denominator. We begin with the following structure:

$$I_5^{\mu\nu} = [I_5^{\mu\nu} - E_{00} g^{\mu\nu}] + E_{00} g^{\mu\nu}. \quad (2.14)$$

The problem of finding an appropriate ansatz for E_{00} has been solved in [16]:

$$E_{00} = \frac{1}{\binom{0}{0}} \sum_{s=1}^5 \binom{0}{s}_5 D_{00}^s, \quad (2.15)$$

where D_{00}^s is the $g^{\mu\nu}$ term of the tensor four point function. The complete result for E_{00} is:

$$E_{00} = -\frac{1}{2} \frac{1}{\binom{0}{0}} \sum_{s=1}^5 \frac{\binom{0}{s}_5}{\binom{s}{5}} \left[\binom{0s}{0s}_5 I_4^s - \sum_{t=1}^5 \binom{0s}{ts}_5 I_3^{st} \right]. \quad (2.16)$$

To demonstrate in details how $()_5$ is cancelled in the square bracket in (2.14), we have to consider four and three point functions separately by analyzing the coefficients of I_4^s and I_3^{st} in (2.9), subtracting the $g^{\mu\nu}$ term according to (2.10) and (2.16). For I_4^s we have

$$\begin{aligned} & \frac{1}{()_5 \binom{0}{0}_5 \binom{s}{s}_5} \\ & \times \left\{ -\binom{0}{j}_5 \binom{0s}{0i}_5 \binom{s}{s}_5 - \binom{s}{j}_5 \binom{0s}{is}_5 \binom{0}{0}_5 + \binom{0}{s}_5 \binom{0s}{0s}_5 \binom{i}{j}_5 \right\}_{ij}^s \\ & \equiv \frac{1}{\binom{0}{0}_5 \binom{s}{s}_5} X_{ij}^s, \end{aligned} \quad (2.17)$$

i.e. we have to show that indeed $()_5$ cancels and we have to give an explicit expression for X_{ij}^s .

A useful property of X_{ij}^s is its symmetry w.r.t. the indices i and j for fixed s . Obviously, the third term in the curly bracket of (2.17) is symmetric since we consider a symmetric determinant. The symmetry of the first two terms means

$$\begin{aligned} & \binom{s}{s}_5 \left[\binom{0}{i}_5 \binom{0j}{0s}_5 - \binom{0}{j}_5 \binom{0i}{0s}_5 \right] \\ & + \binom{0}{0}_5 \left[\binom{s}{i}_5 \binom{0s}{js}_5 - \binom{s}{j}_5 \binom{0s}{is}_5 \right] = 0. \end{aligned} \quad (2.18)$$

The first square bracket of (2.18) can be evaluated using (A.13) of [14], *i.e.*

$$\binom{0}{j}_5 \binom{0i}{0s}_5 = -\binom{0}{0}_5 \binom{0s}{ij}_5 + \binom{0}{i}_5 \binom{0j}{0s}_5, \quad (2.19)$$

and (2.18) then results in:

$$\binom{s}{i}_5 \binom{0s}{js}_5 + \binom{s}{j}_5 \binom{0s}{si}_5 + \binom{s}{s}_5 \binom{0s}{ij}_5 = 0. \quad (2.20)$$

This is proved by multiplication with $()_5$ and using (A.8) of [14] with $r = 2$, *i.e.*

$$\binom{il}{jk}_5 ()_5 = \binom{i}{j}_5 \binom{l}{k}_5 - \binom{i}{k}_5 \binom{l}{j}_5. \quad (2.21)$$

Inserting this, products of three factors of the form $\binom{i}{k}_5$ cancel pairwise, q.e.d.

Further, the following relations (A.11) and (A.12) of [14] are important, *i.e.*

$$\sum_{i=1}^n \binom{0}{i}_5 = ()_5, \quad \sum_{i=1}^n \binom{j}{i}_5 = 0, \quad (j \neq 0). \quad (2.22)$$

As simplest case we now immediately obtain from (2.17) $\{\cdots\}_{ss}^s = 0$, *i.e.* $X_{ss}^s = 0$. Applying (2.22) to (2.17), we see:

$$\sum_{j=1}^5 \{\cdots\}_{ij}^s = -()_5 \binom{0s}{0i}_5 \binom{s}{s}_5, \quad (2.23)$$

and due to the symmetry in i and j we also have:

$$\sum_{i=1}^5 \{\cdots\}_{ij}^s = -()_5 \binom{0s}{0j}_5 \binom{s}{s}_5, \quad (2.24)$$

which gives us a hint of how X_{ij}^s might look, namely due to (2.23) it should contain a term $-\binom{0s}{0i}_5 \binom{0s}{js}_5$. A further contribution, summed over, must vanish. Due to $X_{ss}^s = 0$ it must contain a factor $\binom{0j}{si}_5$. The second factor of this further contribution can only depend on s and has been determined by explicit calculation to be $\binom{0s}{0s}_5$. Thus we conclude

$$X_{ji}^s = X_{ij}^s = -\binom{0s}{0i}_5 \binom{0s}{js}_5 + \binom{0j}{si}_5 \binom{0s}{0s}_5. \quad (2.25)$$

For I_3^{st} a slight generalisation yields the coefficient

$$X_{ij}^{st} = -\binom{0s}{0j}_5 \binom{ts}{is}_5 + \binom{0i}{sj}_5 \binom{ts}{0s}_5, \quad (2.26)$$

with $X_{ij}^s = X_{ij}^{s,t=0}$ and the final result is:

$$I_5^{\mu\nu} = \sum_{i,j=1}^4 q_i^\mu q_j^\nu I_{5,ij} + g^{\mu\nu} E_{00}, \quad (2.27)$$

where

$$I_{5,ij} = \frac{1}{\binom{0}{0}_5} \sum_{s=1}^5 \frac{1}{\binom{s}{s}_5} \left(X_{ij}^{s0} I_4^s - \sum_{t=1}^5 X_{ij}^{st} I_3^{st} \right). \quad (2.28)$$

X_{ij}^{st} is defined in (2.26) and E_{00} in (2.16). I_3^{st} is the scalar three point function obtained by cutting the s and t lines in the five point diagram.

3. Automatization

We have developed a Mathematica package which reduces general tensor five point functions up to rank three. Using `LoopTools` [17] we have performed numerical cross-checks which ensured us about the correctness of the package. We have also made independent checks with sector decomposition and the Mellin–Barnes method. Some examples are given in [18].

In summary, we have developed tools to deal with five point Bhabha amplitudes. The next step is to implement it into the complete numerical calculation.

REFERENCES

- [1] R. Bonciani, A. Ferroglia, P. Mastrolia, E. Remiddi, J.J. van der Bij, *Nucl. Phys.* **B701**, 121 (2004) [[arXiv:hep-ph/0405275](#)]; R. Bonciani, A. Ferroglia, P. Mastrolia, E. Remiddi, J.J. van der Bij, *Nucl. Phys.* **B716**, 280 (2005) [[arXiv:hep-ph/0411321](#)].
- [2] A.A. Penin, *Phys. Rev. Lett.* **95**, 010408 (2005) [[arXiv:hep-ph/0501120](#)].
- [3] S. Actis, M. Czakon, J. Gluza, T. Riemann, *Nucl. Phys.* **B786**, 26 (2007) [[arXiv:0704.2400\[hep-ph\]](#)]; T. Riemann, Fermionic NNLO Corrections to Bhabha Scattering, talk held at this Conference.
- [4] T. Becher, K. Melnikov, *J. High Energy Phys.* **0706**, 084 (2007) [[arXiv:0704.3582\[hep-ph\]](#)].
- [5] M. Melles, *Acta Phys. Pol. B* **28**, 1159 (1997) [[arXiv:hep-ph/9612348](#)].
- [6] S. Jadach, M. Melles, B.F.L. Ward, S.A. Yost, *Phys. Lett.* **B377**, 168 (1996) [[arXiv:hep-ph/9603248](#)].
- [7] S. Jadach, M. Melles, B.F.L. Ward, S.A. Yost, *Phys. Rev.* **D65**, 073030 (2002) [[arXiv:hep-ph/0109279](#)].
- [8] A.B. Arbuzov, D. Haidt, C. Matteuzzi, M. Paganoni, L. Trentadue, *Eur. Phys. J.* **C34**, 267 (2004) [[arXiv:hep-ph/0402211](#)].
- [9] J. Gluza, K. Kajda, T. Riemann, *Comput. Phys. Commun.* **177**, 879 (2007) [[arXiv:0704.2423\[hep-ph\]](#)].
- [10] M. Czakon, *Comput. Phys. Commun.* **175**, 559 (2006) [[arXiv:hep-ph/0511200](#)].
- [11] M. Tentyukov, J. Fleischer, *Nucl. Instrum. Methods*, **A502**, 570 (2003) [[arXiv:hep-ph/0210179](#)].
- [12] J. Fleischer, Application of Mellin–Barnes Representation to the Calculation of Massive Five-Point Functions in Bhabha Scattering, talk given at the Conference on Frontiers in Perturbative Quantum Field Theory, Bielefeld, Germany, 2007.
- [13] G. Passarino, M.J.G. Veltman, *Nucl. Phys.* **B160**, 151 (1979).
- [14] D.B. Melrose, *Nuovo Cim.* **40**, 181 (1965).

- [15] J. Fleischer, F. Jegerlehner, O.V. Tarasov, *Nucl. Phys.* **B566**, 423 (2000) [arXiv:hep-ph/9907327].
- [16] A. Denner, S. Dittmaier, *Nucl. Phys.* **B658**, 175 (2003) [arXiv:hep-ph/0212259]; A. Denner, S. Dittmaier, *Nucl. Phys.* **B734** (2006) [arXiv:hep-ph/0509141].
- [17] T. Hahn, M. Perez-Victoria, *Comput. Phys. Commun.* **118**, 153 (1999) [arXiv:hep-ph/9807565].
- [18] Transparencies of this talk, <http://www.us.edu.pl/~us2007>